

93 Analytical solution of differential equations

1. Nonlinear differential equation

93.1. (Separable differential equation. See AMBS Ch 38–39.) Find analytical solution formulas for the following initial value problems. In each case sketch the graphs of the solutions and determine the half-life. See: P. Atkins and L. Jones, *Chemical Principles. The Quest for Insight*. Freeman, New York, second edition, 2002, pp. 698–706.

(a) First order rate law:

$$\begin{aligned}u' &= -ku, \quad t > 0, \\u(0) &= u_0.\end{aligned}$$

(b) Second order rate law:

$$\begin{aligned}u' &= -ku^2, \quad t > 0, \\u(0) &= u_0.\end{aligned}$$

(c) Third order rate law:

$$\begin{aligned}u' &= -ku^3, \quad t > 0, \\u(0) &= u_0.\end{aligned}$$

2. Linear differential equation

2.1 Linear differential equation—first order

$$(93.1) \quad u' + a(t)u = f(t).$$

Here $u = u(t)$ is an unknown function of an independent variable t . The equation is called *homogeneous* if $f(t) \equiv 0$ and *nonhomogeneous* otherwise. The differential operator $Lu = u' + a(t)u$ has *constant coefficient* if $a(t) = a$ is constant and it has *variable coefficient* otherwise. The equation is said to be a *linear equation*, because the operator L is a *linear operator*:

$$L(\alpha u + \beta v) = \alpha Lu + \beta Lv, \quad (\alpha, \beta \in \mathbf{R}, \quad u = u(t), \quad v = v(t))$$

i.e., it preserves linear combinations of functions. Check this!

Solution method: *integrating factor*. See AMBS Ch 35.1–2.

93.2. (constant coefficient, homogeneous) Solve the following. Sketch the graph of the solution.

(a)

$$\begin{aligned}u' + 2u &= 0, \quad t > 0, \\u(0) &= u_0.\end{aligned} \quad \text{Solution: } u(t) = e^{-2t}u_0.$$

(b)

$$\begin{aligned}u' - 2u &= 0, \quad t > 0, \\u(0) &= u_0.\end{aligned} \quad \text{Solution: } u(t) = e^{2t}u_0.$$

93.3. (constant coefficient, nonhomogeneous) Solve the following.

(a)

$$\begin{aligned}u' + 2u &= f(t), \quad t > 0, \\u(0) &= u_0.\end{aligned} \quad \text{Solution: } u(t) = e^{-2t}u_0 + \int_0^t e^{-2(t-s)}f(s)ds.$$

(b)

$$\begin{aligned} u' - 2u &= f(t), \quad t > 0, \\ u(0) &= u_0. \end{aligned} \quad \text{Solution: } u(t) = e^{2t}u_0 + \int_0^t e^{2(t-s)}f(s) \, ds.$$

93.4. (constant coefficient, nonhomogeneous) Solve the following.

$$\begin{aligned} u' + au &= f(t), \quad t > 0, \\ u(0) &= u_0 \end{aligned} \quad \text{Solution: } u(t) = e^{-at}u_0 + \int_0^t e^{-a(t-s)}f(s) \, ds.$$

93.5. (variable coefficient, nonhomogeneous) Solve the following.

$$\begin{aligned} u' + 2tu &= f(t), \quad t > 0, \\ u(0) &= u_0. \end{aligned} \quad \text{Solution: } u(t) = e^{-t^2}u_0 + \int_0^t e^{-(t^2-s^2)}f(s) \, ds.$$

2.2 Linear differential equation—second order—constant coefficients

$$(93.2) \quad u'' + a_1u' + a_0u = f(t).$$

The equation is called *homogeneous* if $f(t) \equiv 0$ and *nonhomogeneous* otherwise. We assume that the differential operator $Lu = u'' + a_1u' + a_0u$ has *constant coefficients* a_1 and a_0 . Check that the operator L is linear!

Variable coefficients: Linear differential equations of second order with variable coefficients $u'' + a_1(t)u' + a_0(t)u = f(t)$, cannot be solved analytically, except in some special cases. One such case can be found in AMBS Ch 35.6. We do not discuss this here.

Homogeneous equation

See AMBS Ch 35.3–35.4. The homogeneous equation (93.2) may be written

$$(93.3) \quad D^2u + a_1Du + a_0u = 0,$$

or

$$P(D)u = 0,$$

where

$$P(r) = r^2 + a_1r + a_0$$

is the *characteristic polynomial* of the equation. The *characteristic equation* $P(r) = 0$ has two roots r_1 and r_2 . Hence $P(r) = (r - r_1)(r - r_2)$. All solutions of equation (93.2) are obtained as linear combinations

$$(93.4) \quad \begin{aligned} u(t) &= c_1e^{r_1t} + c_2e^{r_2t}, & \text{if } r_1 \neq r_2, \\ u(t) &= (c_1 + c_2t)e^{r_1t}, & \text{if } r_1 = r_2, \end{aligned}$$

where c_1, c_2 are arbitrary coefficients. The coefficients may be determined from an initial condition of the form

$$u(0) = u_0, \quad u'(0) = u_1.$$

The formula (93.4) is called the *general solution* of homogeneous linear equation (93.3).

Example 93.1. We solve

$$u'' + u' - 12u = 0; \quad u(0) = u_0, \quad u'(0) = u_1.$$

The equation is written $(D^2 + D - 12)u = 0$ and the characteristic equation is $r^2 + r - 12 = 0$ with roots $r_1 = 3, r_2 = -4$. The general solution is

$$u(t) = c_1e^{3t} + c_2e^{-4t}$$

with the derivative

$$u'(t) = 3c_1e^{3t} - 4c_2e^{-4t}.$$

The initial condition gives

$$\begin{aligned} u_0 &= u(0) = c_1 + c_2 \\ u_1 &= u'(0) = 3c_1 - 4c_2 \end{aligned}$$

which implies $c_1 = (4u_0 + u_1)/7$, $c_2 = (3u_0 - u_1)/7$. The solution is

$$u(t) = \frac{4u_0 + u_1}{7}e^{3t} + \frac{3u_0 - u_1}{7}e^{-4t}.$$

93.6. Prove the solution formula (93.4) by writing the equation as

$$P(D)u = (D - r_1)(D - r_2)u = 0$$

and by solving two first order equations $(D - r_1)v = 0$ and $(D - r_2)u = v$ as in Problems 93.2 and 93.3.

93.7. Write the following equations as $P(D)u = 0$ and solve the initial value problem. Choose numerical values for the constants and sketch the graph of the solution. Solve the problem with your MATLAB program `my_ode.m`.

(a) $u'' - u' - 2u = 0$; $u(0) = u_0$, $u'(0) = u_1$.

(b) $u'' - k^2u = 0$; $u(0) = u_0$, $u'(0) = u_1$.

(c) $u'' + 4u' + 4u = 0$; $u(0) = u_0$, $u'(0) = u_1$.

93.8. Solve the *boundary value problem*

$$\begin{aligned} u''(x) - k^2u(x) &= 0, \quad 0 < x < L, \\ u(0) &= 0, \quad u(L) = u_L. \end{aligned}$$

Complex roots

If the characteristic polynomial $P(r)$ has real coefficients, then its roots are real or a complex conjugate pair. In the latter case we have $r_1 = \alpha + i\omega$ and $r_2 = \alpha - i\omega$ and the solution (93.4) becomes (see AMBS Ch 33.2 for the definition of $\exp(z)$ with a complex variable z)

$$\begin{aligned} u(t) &= c_1e^{(\alpha+i\omega)t} + c_2e^{(\alpha-i\omega)t} \\ &= e^{\alpha t} \left(c_1e^{i\omega t} + c_2e^{-i\omega t} \right) \\ &= e^{\alpha t} \left(c_1(\cos(\omega t) + i\sin(\omega t)) + c_2(\cos(\omega t) - i\sin(\omega t)) \right) \\ &= e^{\alpha t} \left(d_1\cos(\omega t) + d_2\sin(\omega t) \right), \end{aligned}$$

with $d_1 = c_1 + c_2$, $d_2 = i(c_1 - c_2)$.

93.9. Write the equation as $P(D)u = 0$ and solve the initial value problem. Choose numerical values for the constants and sketch the graph of the solution. Solve the problem with your MATLAB program `my_ode.m`.

(a) $u'' + 4u' + 13u = 0$; $u(0) = u_0$, $u'(0) = u_1$.

(b) $u'' + \omega^2u = 0$; $u(0) = u_0$, $u'(0) = u_1$.

Nonhomogeneous equation

See AMBS Ch 35.5. The solution of the nonhomogeneous equation $P(D)u = f(t)$ is given by

$$(93.5) \quad u(t) = u_h(t) + u_p(t),$$

where u_h is the general solution (93.4) of the corresponding homogeneous equation, i.e., $P(D)u_h = 0$, and u_p is a *particular solution* of the nonhomogeneous equation, i.e., $P(D)u_p = f(t)$. Prove this!

A particular solution can sometimes be found by guess-work: we make an Ansatz for u_p of the same form as f .

Example 93.2. $u'' - u' - 2u = t$. Here $f(t) = t$ is a polynomial of degree 1 and we make the Ansatz $u_p(t) = At + B$, i.e., a polynomial of degree 1. Substitution into the equation gives $-A - 2(At + B) = t$. Identification of coefficients gives $A = -\frac{1}{2}$, $B = \frac{1}{4}$, so that $u_p(t) = \frac{1}{4} - \frac{1}{2}t$. The general solution of the homogeneous equation is $u_h(t) = c_1e^{-t} + c_2e^{2t}$, see Problem 93.7 (a). Hence we get

$$u(t) = u_h(t) + u_p(t) = c_1e^{-t} + c_2e^{2t} + \frac{1}{4} - \frac{1}{2}t.$$

93.10. Solve the following.

- (a) $u'' - u' - 2u = e^t$ Ansatz: $u_p(t) = Ae^t$
- (b) $u'' - u' - 2u = \cos(t)$ Ansatz: $u_p(t) = A\cos(t) + B\sin(t)$
- (c) $u'' - u' - 2u = t^3$ Ansatz: $u_p(t) = At^3 + Bt^2 + Ct + D$
- (d) $u'' - u' - 2u = e^{-t}$ Ansatz: $u_p(t) = Ate^{-t}$

Re-writing as a system of first order equations

By setting $w_1 = u$, $w_2 = u'$, $w = \begin{bmatrix} w_1 \\ w_2 \end{bmatrix}$, we can re-write (93.2) as a system of first order equations

$$w'(t) = Aw(t) + F(t); \quad w(0) = w_0,$$

where

$$w_0 = \begin{bmatrix} u_0 \\ u_1 \end{bmatrix}, \quad A = \begin{bmatrix} 0 & 1 \\ -a_0 & -a_1 \end{bmatrix}, \quad F(t) = \begin{bmatrix} 0 \\ f(t) \end{bmatrix}.$$

To see this we compute

$$w' = \begin{bmatrix} u' \\ u'' \end{bmatrix} = \begin{bmatrix} u' \\ -a_0u - a_1u' + f(t) \end{bmatrix} = \begin{bmatrix} w_2 \\ -a_0w_1 - a_1w_2 + f(t) \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -a_0 & -a_1 \end{bmatrix} \begin{bmatrix} w_1 \\ w_2 \end{bmatrix} + \begin{bmatrix} 0 \\ f(t) \end{bmatrix}.$$

2.3 System of linear differential equations of first order

Constant coefficients—homogeneous equations

We now consider

$$(93.6) \quad \begin{aligned} u' + Au &= 0, \quad t > 0, \\ u(0) &= u_0, \end{aligned}$$

where $u(t), u_0 \in \mathbf{R}^d$, and $A \in \mathbf{R}^{d \times d}$ is a constant matrix of coefficients. We assume that the matrix A is diagonalizable. This means that there is a diagonal matrix $D = \text{diag}(\lambda_1, \dots, \lambda_d)$ of eigenvalues and a matrix $P = [g_1, \dots, g_d]$ of eigenvectors such that P is invertible and

$$AP = PD, \quad A = PDP^{-1}, \quad P^{-1}AP = D.$$

We define

$$e^{-tD} = \text{diag}(e^{-t\lambda_1}, \dots, e^{-t\lambda_d})$$

and

$$e^{-tA} = Pe^{-tD}P^{-1}.$$

It is now easy to check that the solution of (93.6) is given by

$$(93.7) \quad u(t) = e^{-tA}u_0.$$

Compare this with the scalar case in Problem 93.4. The solution (93.7) may also be written as a linear combination

$$u(t) = c_1 e^{-t\lambda_1} g_1 + \dots + c_d e^{-t\lambda_d} g_d.$$

The coefficients are determined by the initial condition:

$$u_0 = u(0) = c_1 g_1 + \dots + c_d g_d.$$

This is a system of linear equations for the coefficients.

The case when A is not diagonalizable is more complicated, but can also be handled by eigenvalue techniques. We do not discuss this here.

93.11. Solve the system

$$u' + Au = 0; \quad u(0) = u_0, \quad A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}, \quad u_0 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}.$$

93.12. Write the equations in Problems 93.7 and 93.9 as systems of first order equations and solve by the “eigenvector method”.

System of nonhomogeneous equations

The solution of the nonhomogeneous system

$$u' + Au = f(t); \quad u(0) = u_0,$$

is given by

$$u(t) = e^{-At}u_0 + \int_0^t e^{-A(t-s)} f(s) ds.$$

This is proved in the same way as Problem 93.4. The integrating factor is e^{At} .

Variable coefficients

Linear systems of differential equations with variable coefficients

$$u' + A(t)u = f(t); \quad u(0) = u_0,$$

cannot be solved analytically, except in some special cases.

Answers and solutions

93.1. Reaction of order 1 (decay rate of order 1):

$$\begin{cases} u' = -ku \\ u(0) = u_0 \end{cases}$$

$$u(t) = u_0 e^{-kt}$$

The half-life $T_{1/2}$ is given by

$$u(T_{1/2}) = u_0 e^{-kT_{1/2}} = \frac{1}{2}u_0,$$

which leads to

$$T_{1/2} = \frac{\log(2)}{k}.$$

Reaction of order $n > 1$ (decay rate of order $n > 1$):

$$\begin{aligned} & \begin{cases} u' = -ku^n \\ u(0) = u_0 \end{cases} \\ & \frac{du}{u^n} = -k dt \\ & \int_{u_0}^{u(T)} u^{-n} du = - \int_0^T k dt \\ & \left[\frac{u^{-n+1}}{-n+1} \right]_{u_0}^{u(T)} = -kT \\ & u(T)^{-n+1} - u_0^{-n+1} = (n-1)kT \\ & \frac{1}{u(T)^{n-1}} = \frac{1}{u_0^{n-1}} + (n-1)kT = \frac{1 + (n-1)u_0^{n-1}kT}{u_0^{n-1}} \\ & u(T) = \frac{u_0}{(1 + (n-1)u_0^{n-1}kT)^{1/(n-1)}} \end{aligned}$$

The half-life $T_{1/2}$ is given by

$$u(T_{1/2}) = \frac{u_0}{(1 + (n-1)u_0^{n-1}kT_{1/2})^{1/(n-1)}} = \frac{1}{2}u_0$$

which implies

$$T_{1/2} = \frac{2^{n-1} - 1}{(n-1)u_0^{n-1}k}$$

93.7.

- (a) $u(t) = \frac{1}{3}(2u_0 - u_1)e^{-t} + \frac{1}{3}(u_0 + u_1)e^{2t}$.
- (b) $u(t) = c_1 e^{kt} + c_2 e^{-kt} = d_1 \cosh(kt) + d_2 \sinh(kt)$, $d_1 = c_1 + c_2$, $d_2 = c_1 - c_2$. The initial condition gives $u(t) = \frac{1}{2}(u_0 + u_1/k)e^{kt} + \frac{1}{2}(u_0 - u_1/k)e^{-kt}$ or alternatively $u(t) = u_0 \cosh(kt) + (u_1/k) \sinh(kt)$.
- (c) $u(t) = (u_0 + (2u_0 + u_1)t)e^{-2t}$.

93.8. $u(x) = u_L \sinh(kx) / \sinh(kL)$.

93.9.

- (a) $u(t) = e^{-2t}(u_0 \cos(3t) + \frac{1}{3}(2u_0 + u_1) \sin(3t))$.
- (b) $u(t) = u_0 \cos(\omega t) + (u_1/\omega) \sin(\omega t)$. Compare to Problem 93.7 (b).

93.10.

- (a) $u(t) = c_1 e^{-t} + c_2 e^{2t} - \frac{1}{2}e^t$.

(b) $u(t) = c_1 e^{-t} + c_2 e^{2t} - \frac{3}{10} \cos(t) - \frac{1}{10} \sin(t).$

(c) $u(t) = c_1 e^{-t} + c_2 e^{2t} - \frac{1}{2} t^3 + \frac{3}{4} t^2 - \frac{9}{4} t + \frac{15}{8}.$

(d) $u(t) = c_1 e^{-t} + c_2 e^{2t} - \frac{1}{3} t e^{-t}.$ Note: the Ansatz $u_p(t) = A e^{-t}$ does not work, because e^{-t} is a solution of the homogeneous equation, $P(D)e^{-t} = 0$, i.e., e^{-t} is contained in u_h .

93.11.

$$D = \begin{bmatrix} 3 & 0 \\ 0 & -1 \end{bmatrix}, \quad P = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}, \quad P^T = P^{-1}$$

$$e^{-tA} = P e^{-tD} P^T = \frac{1}{2} \begin{bmatrix} e^{-3t} + e^t & e^{-3t} - e^t \\ e^{-3t} - e^t & e^{-3t} + e^t \end{bmatrix}.$$

$$u(t) = e^{-tA} u_0 = \frac{1}{2} \begin{bmatrix} e^{-3t} + e^t & e^{-3t} - e^t \\ e^{-3t} - e^t & e^{-3t} + e^t \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 3e^{-3t} - e^t \\ 3e^{-3t} + e^t \end{bmatrix}.$$

Alternatively

$$\begin{aligned} u(t) &= c_1 e^{-t\lambda_1} g_1 + c_2 e^{-t\lambda_2} g_2 = c_1 e^{-3t} \begin{bmatrix} 1 \\ 1 \end{bmatrix} + c_2 e^t \begin{bmatrix} 1 \\ -1 \end{bmatrix} \\ &= \frac{3}{2} e^{-3t} \begin{bmatrix} 1 \\ 1 \end{bmatrix} - \frac{1}{2} e^t \begin{bmatrix} 1 \\ -1 \end{bmatrix}. \end{aligned}$$

2002-11-22 /stig